The Neck Disability Index

Pa	tient name:		File#	Date:
Please read instructions: This questionnaire has been designed to give the doctor information as to how your neck pain has affected your ability to manage everyday life. Please answer every section and mark in each section only the ONE box that applies to you. We realize that you may consider that two of the statements in any one section relate to you, but please just mark the box that most closely describes your problem.				
SECTION 1-PAIN INTENSITY		SECTION 6-CONCENTRATION		
	I have no pain at the moment. The pain is very mild at the moment. The pain is moderate at the moment. The pain is fairly severe at the moment. The pain is very severe at the moment. The pain is the worst imaginable at the moment.		I can concentrate fully when I want I can concentrate fully when I wan I have a fair degree of difficulty in I have a lot of difficulty in concentr I have a great deal of difficulty in c I cannot concentrate at all.	t to, with slight difficulty. concentrating when I want to. ating when I want to.
SECTION 2-PERSONAL CARE (Washing, Dressing, etc.)		SECTION 7-WORK		
	I can look after myself normally, without causing extra pain. I can look after myself normally, but it causes extra pain. It is painful to look after myself and I am slow and careful. I need some help, but manage most of my personal care. I need help every day in most aspects of self care. I do not get dressed; I wash with difficulty and stay in bed.		I can do as much work as I want to. I can do my usual work, but no mor I can do most of my usual work, but cannot do my usual work. I can hardly do any work at all. I can't do any work at all.	e.
SECTION 3-LIFTING		SECTION 8-DRIVING		
	I can lift heavy weights without extra pain. I can lift heavy weights, but it gives extra pain. Pain prevents me from lifting heavy weights off the floor, but I can manage if they are conveniently positioned, for example, on a table. Pain prevents me from lifting heavy weights off the floor, but I can manage light to medium weights if they are conveniently positioned. I can lift very light weights. I cannot lift or carry anything at all.		I can drive my car without any neck I can drive my car as long as I want I can drive my car as long as I want neck. I can't drive my car as long as I wan in my neck. I can hardly drive at all, because of I can't drive my car at all.	t, with slight pain in my neck. The with moderate pain in my at, because of moderate pain
SECTION 4-READING		SECTION 9-SLEEPING		
	I can read as much as I want to, with no pain in my neck. I can read as much as I want to, with slight pain in my neck. I can read as much as I want to, with moderate pain in my neck. I can't read as much as I want, because of moderate pain in my neck. I can hardly read at all, because of severe pain in my neck. I cannot read at all.		I have no trouble sleeping. My sleep is slightly disturbed (less to My sleep is mildly disturbed (1-2 h My sleep is moderately disturbed (3-5 h My sleep is greatly disturbed (3-5 h My sleep is completely di	rs sleepless). 2-3 hrs sleepless). rs sleepless).
SECTION 5-HEADACHES		SECTION 10-RECREATION		
	I have no headaches at all. I have slight headaches that come infrequently. I have moderate headaches that come infrequently. I have moderate headaches that come frequently. I have severe headaches that come frequently. I have headaches almost all the time.		I am able to engage in all my recre- pain at all. I am able to engage in all my recre- neck pain at all. I am able to engage in most, but no activities, because of pain in my ne- I am able to engage in few of my re- pain in my neck. I can hardly do any recreation activ- neck. I can't do any recreation activities a	ation activities, with some t all, of my usual recreation ck. ccreation activities, because of ities, because of pain in my

Instructions:

- 1. The NDI is scored in the same way as the Oswestry Disability Index.
- 2. Using this system, a score of 10-28% (i.e., 5-14 points) is considered by the authors to constitute mild disability; 30-48% is moderate; 50-68% is severe; 72% or more is complete.